

Introductory
STATISTICS

9TH EDITION



Neil
WEISS

Chapter 4

Probability Concepts



Section 4.1

Probability Basics



Definition 4.1

Probability for Equally Likely Outcomes (f/N Rule)

Suppose an experiment has N possible outcomes, all equally likely. An event that can occur in f ways has probability f/N of occurring:

$$\text{Probability of an event} = \frac{f}{N}.$$

Number of ways event can occur

Total number of possible outcomes


The diagram shows the equation 'Probability of an event = f/N.' with two arrows. One arrow points from the text 'Number of ways event can occur' to the variable 'f' in the numerator. The other arrow points from the text 'Total number of possible outcomes' to the variable 'N' in the denominator.

Figure 4.1

Possible outcomes for rolling a pair of dice






Figure 4.2

Two computer simulations of tossing a balanced coin 100 times



Key Fact 4.1

Basic Properties of Probabilities

Property 1: The probability of an event is always between 0 and 1, inclusive.

Property 2: The probability of an event that cannot occur is 0. (An event that cannot occur is called an **impossible event**.)

Property 3: The probability of an event that must occur is 1. (An event that must occur is called a **certain event**.)

Section 4.2

Events



Definition 4.2


Sample Space and Event

Sample space: The collection of all possible outcomes for an experiment.


Event: A collection of outcomes for the experiment, that is, any subset of the sample space. An event **occurs** if and only if the outcome of the experiment is a member of the event.

Figure 4.9


Venn diagrams for (a) event (not E), (b) event ($A \& B$), and (c) event (A or B)



(a)



(b)



(c)

Definition 4.3

Relationships Among Events

(not E): The event “ E does not occur”

(A & B): The event “both A and B occur”

(A or B): The event “either A or B or both occur”

Definition 4.4

Mutually Exclusive Events

Two or more events are **mutually exclusive events** if no two of them have outcomes in common.

Figure 4.14

- (a) Two mutually exclusive events;
- (b) two non-mutually exclusive events






Figure 4.15


- (a) Three mutually exclusive events;
- (b) three non-mutually exclusive events;
- (c) three non-mutually exclusive events



(a)



(b)



(c)

Section 4.3

Some Rules of Probability



Formula 4.1

The Special Addition Rule

If event A and event B are mutually exclusive, then

$$P(A \text{ or } B) = P(A) + P(B).$$

More generally, if events A, B, C, \dots are mutually exclusive, then

$$P(A \text{ or } B \text{ or } C \text{ or } \dots) = P(A) + P(B) + P(C) + \dots.$$

Formula 4.2

The Complementation Rule

For any event E ,

$$P(E) = 1 - P(\text{not } E).$$

Formula 4.3

The General Addition Rule

If A and B are any two events, then

$$P(A \text{ or } B) = P(A) + P(B) - P(A \& B).$$

Section 4.4

Contingency Tables; Joint and Marginal Probabilities



Table 4.6 Contingency table for age and rank of faculty members

		Rank				Total
		Full professor R_1	Associate professor R_2	Assistant professor R_3	Instructor R_4	
Age (yr)	Under 30 A_1	2	3	57	6	68
	30–39 A_2	52	170	163	17	402
	40–49 A_3	156	125	61	6	348
	50–59 A_4	145	68	36	4	253
	60 & over A_5	75	15	3	0	93
	Total	430	381	320	33	1164

Table 4.7 Joint probability distribution corresponding to Table 4.6

		Rank				$P(A_i)$
		Full professor R_1	Associate professor R_2	Assistant professor R_3	Instructor R_4	
Age (yr)	Under 30 A_1	0.002	0.003	0.049	0.005	0.058
	30–39 A_2	0.045	0.146	0.140	0.015	0.345
	40–49 A_3	0.134	0.107	0.052	0.005	0.299
	50–59 A_4	0.125	0.058	0.031	0.003	0.217
	60 & over A_5	0.064	0.013	0.003	0.000	0.080
	$P(R_j)$	0.369	0.327	0.275	0.028	1.000

Section 4.5

Conditional Probability



Definition 4.6

Conditional Probability

The probability that event B occurs given that event A occurs is called a **conditional probability**. It is denoted $P(B | A)$, which is read “the probability of B given A .” We call A the **given event**.

Formula 4.4

The Conditional Probability Rule

If A and B are any two events with $P(A) > 0$, then

$$P(B | A) = \frac{P(A \& B)}{P(A)}.$$

Table 4.9

Joint probability distribution of marital status and gender

		Marital status				$P(S_i)$
		Single M_1	Married M_2	Widowed M_3	Divorced M_4	
Gender	Male S_1	0.138	0.290	0.012	0.044	0.484
	Female S_2	0.114	0.291	0.051	0.060	0.516
	$P(M_j)$	0.252	0.581	0.063	0.104	1.000

Section 4.6

The Multiplication Rule; Independence



Formula 4.5


The General Multiplication Rule

If A and B are any two events, then

$$P(A \& B) = P(A) \cdot P(B | A).$$

Figure 4.25

Tree diagram for student-selection problem



Definition 4.7

Independent Events

Event B is said to be **independent** of event A if $P(B | A) = P(B)$.

Formula 4.7

The Special Multiplication Rule

If events A, B, C, \dots are independent, then

$$P(A \& B \& C \& \dots) = P(A) \cdot P(B) \cdot P(C) \dots$$

Section 4.7

Bayes's Rule



Table 4.11 & 4.12

Percentage distribution for region of residence and percentage of seniors in each region


Region	Percentage of U.S. population	Percentage seniors
Northeast	18.3	13.6
Midwest	22.2	12.8
South	36.3	12.5
West	23.2	11.2
	100.0	

Probabilities derived from Table 4.11

$$\begin{aligned}P(R_1) &= 0.183 & P(S | R_1) &= 0.136 \\P(R_2) &= 0.222 & P(S | R_2) &= 0.128 \\P(R_3) &= 0.363 & P(S | R_3) &= 0.125 \\P(R_4) &= 0.232 & P(S | R_4) &= 0.112\end{aligned}$$

Figure 4.27

Tree diagram for calculating $P(S)$, using the rule of total probability



Section 4.8

Counting Rules



Definition 4.8

Factorials

The product of the first k positive integers (counting numbers) is called **k factorial** and is denoted **$k!$** . In symbols,

$$k! = k(k - 1) \cdots 2 \cdot 1.$$

We also define $0! = 1$.

Table 4.14

Possible permutations of three letters from the collection of five letters

<i>abc</i>	<i>abd</i>	<i>abe</i>	<i>acd</i>	<i>ace</i>	<i>ade</i>	<i>bcd</i>	<i>bce</i>	<i>bde</i>	<i>cde</i>
<i>acb</i>	<i>adb</i>	<i>aeb</i>	<i>adc</i>	<i>aec</i>	<i>aed</i>	<i>bdc</i>	<i>bec</i>	<i>bed</i>	<i>ced</i>
<i>bac</i>	<i>bad</i>	<i>bae</i>	<i>cad</i>	<i>cae</i>	<i>dae</i>	<i>cbd</i>	<i>cbe</i>	<i>dbe</i>	<i>dce</i>
<i>bca</i>	<i>bda</i>	<i>bea</i>	<i>cda</i>	<i>cea</i>	<i>dea</i>	<i>cdb</i>	<i>ceb</i>	<i>deb</i>	<i>dec</i>
<i>cab</i>	<i>dab</i>	<i>eab</i>	<i>dac</i>	<i>eac</i>	<i>ead</i>	<i>dbc</i>	<i>ebc</i>	<i>ebd</i>	<i>ecd</i>
<i>cba</i>	<i>dba</i>	<i>eba</i>	<i>dca</i>	<i>eca</i>	<i>eda</i>	<i>dcb</i>	<i>ecb</i>	<i>edb</i>	<i>edc</i>

Formula 4.10

The Permutations Rule

The number of possible permutations of r objects from a collection of m objects is given by the formula

$${}_m P_r = \frac{m!}{(m-r)!}.$$

Figure 4.29

Calculating the number of outcomes in which exactly 2 of the 5 TVs selected are defective

