

Introductory
STATISTICS

9TH EDITION



Neil
WEISS

Chapter 7

The Sampling Distribution of the Sample Mean



Section 7.1

Sampling Error; the Need for Sampling Distributions



Definition 7.1

Sampling Error

Sampling error is the error resulting from using a sample to estimate a population characteristic.

Definition 7.2

Sampling Distribution of the Sample Mean

For a variable x and a given sample size, the distribution of the variable is called the **sampling distribution of the sample mean.**

Table 7.2

Possible samples
and sample means
for samples of size 2

Sample	Heights	\bar{x}
A, B	76, 78	77.0
A, C	76, 79	77.5
A, D	76, 81	78.5
A, E	76, 86	81.0
B, C	78, 79	78.5
B, D	78, 81	79.5
B, E	78, 86	82.0
C, D	79, 81	80.0
C, E	79, 86	82.5
D, E	81, 86	83.5

Figure 7.1

Dotplot for the sampling distribution of the sample mean for samples of size 2 ($n = 2$)




Figure 7.3

Dotplots for the sampling distributions of the sample mean for the heights of the five starting players for samples of sizes 1, 2, 3, 4, and 5




Table 7.4

Sample size and sampling error illustrations for the heights of the basketball players (“No.” is an abbreviation of “Number”)

Sample size <i>n</i>	No. possible samples	No. within 1'' of μ	% within 1'' of μ	No. within 0.5'' of μ	% within 0.5'' of μ
1	5	2	40%	0	0%
2	10	3	30%	2	20%
3	10	5	50%	2	20%
4	5	4	80%	3	60%
5	1	1	100%	1	100%

Section 7.2

The Mean and Standard Deviation of the Sample Mean



Formula 7.1

Mean of the Sample Mean

For samples of size n , the mean of the variable \bar{x} equals the mean of the variable under consideration. In symbols,

$$\mu_{\bar{x}} = \mu.$$

Formula 7.2

Standard Deviation of the Sample Mean

For samples of size n , the standard deviation of the variable \bar{x} equals the standard deviation of the variable under consideration divided by the square root of the sample size. In symbols,

$$\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}.$$

Section 7.3

The Sampling Distribution of the Sample Mean



Output 7.1

Histogram of the sample means for 1000 samples of four IQs with superimposed normal curve




Figure 7.4

(a) Normal distribution for IQs; (b) sampling distribution of the sample mean for $n = 4$; (c) sampling distribution of the sample mean for $n = 16$






Figure 7.5

Relative-frequency histogram for household size



Output 7.2

Histogram of the sample means for 1000 samples of 30 household sizes with superimposed normal curve



Key Fact 7.4

Sampling Distribution of the Sample Mean

Suppose that a variable x of a population has mean μ and standard deviation σ . Then, for samples of size n ,

- the mean of \bar{x} equals the population mean, or $\mu_{\bar{x}} = \mu$;
- the standard deviation of \bar{x} equals the population standard deviation divided by the square root of the sample size, or $\sigma_{\bar{x}} = \sigma/\sqrt{n}$;
- if x is normally distributed, so is \bar{x} , regardless of sample size; and
- if the sample size is large, \bar{x} is approximately normally distributed, regardless of the distribution of x .

Figure 7.6

Sampling distributions of the sample mean for (a) normal, (b) reverse-J-shaped, and (c) uniform variables

