

Introductory
STATISTICS

9TH EDITION



Neil
WEISS

Chapter 11

Inferences for Population Standard Deviations




Section 11.1

Inferences for One Population Standard Deviation



Figure 11.1

χ^2 -curves for $df = 5, 10,$ and 19



Key Fact 11.2

The Sampling Distribution of the Sample Standard Deviation[†]

Suppose that a variable of a population is normally distributed with standard deviation σ . Then, for samples of size n , the variable

$$\chi^2 = \frac{n-1}{\sigma^2} s^2$$

has the chi-square distribution with $n - 1$ degrees of freedom.

Procedure 11.1

One-Standard-Deviation χ^2 -Test

Purpose To perform a hypothesis test for a population standard deviation, σ

Assumptions

1. Simple random sample
2. Normal population

Step 1 The null hypothesis is $H_0: \sigma = \sigma_0$, and the alternative hypothesis is

$$\begin{array}{ccc} H_a: \sigma \neq \sigma_0 & \text{or} & H_a: \sigma < \sigma_0 & \text{or} & H_a: \sigma > \sigma_0 \\ \text{(Two tailed)} & & \text{(Left tailed)} & & \text{(Right tailed)} \end{array}$$

Step 2 Decide on the significance level, α .

Step 3 Compute the value of the test statistic

$$\chi^2 = \frac{n - 1}{\sigma_0^2} s^2$$

and denote that value χ_0^2 .


Procedure 11.1 (cont.)

CRITICAL-VALUE APPROACH

Step 4 The critical value(s) are

$\chi^2_{1-\alpha/2}$ and $\chi^2_{\alpha/2}$ (Two tailed) or $\chi^2_{1-\alpha}$ (Left tailed) or χ^2_{α} (Right tailed)

with $df = n - 1$. Use Table VII to find the critical value(s).




Step 5 If the value of the test statistic falls in the rejection region, reject H_0 ; otherwise, do not reject H_0 .

OR

P-VALUE APPROACH

Step 4 The χ^2 -statistic has $df = n - 1$. Obtain the P -value by using technology.



Step 5 If $P \leq \alpha$, reject H_0 ; otherwise, do not reject H_0 .

Step 6 Interpret the results of the hypothesis test.

Procedure 11.2

One-Standard-Deviation χ^2 -Interval Procedure

Purpose To find a confidence interval for a population standard deviation, σ

Assumptions

1. Simple random sample
2. Normal population

Step 1 For a confidence level of $1 - \alpha$, use Table VII to find $\chi_{1-\alpha/2}^2$ and $\chi_{\alpha/2}^2$ with $df = n - 1$.

Step 2 The confidence interval for σ is from

$$\sqrt{\frac{n-1}{\chi_{\alpha/2}^2}} \cdot s \quad \text{to} \quad \sqrt{\frac{n-1}{\chi_{1-\alpha/2}^2}} \cdot s,$$

where $\chi_{1-\alpha/2}^2$ and $\chi_{\alpha/2}^2$ are found in Step 1, n is the sample size, and s is computed from the sample data obtained.

Step 3 Interpret the confidence interval.


Section 11.2

Inferences for Two Population Standard Deviations, Using Independent Samples



Figure 11.7

Two different F-curves



Key Fact 11.5

Distribution of the F -Statistic for Comparing Two Population Standard Deviations

Suppose that the variable under consideration is normally distributed on each of two populations. Then, for independent samples of sizes n_1 and n_2 from the two populations, the variable

$$F = \frac{s_1^2/\sigma_1^2}{s_2^2/\sigma_2^2}$$

has the F -distribution with $df = (n_1 - 1, n_2 - 1)$.

Procedure 11.3

Two-Standard-Deviations *F*-Test

Purpose To perform a hypothesis test to compare two population standard deviations, σ_1 and σ_2

Assumptions

1. Simple random samples
2. Independent samples
3. Normal populations

Step 1 The null hypothesis is $H_0: \sigma_1 = \sigma_2$, and the alternative hypothesis is

$$\begin{array}{ccc} H_a: \sigma_1 \neq \sigma_2 & \text{or} & H_a: \sigma_1 < \sigma_2 & \text{or} & H_a: \sigma_1 > \sigma_2 \\ \text{(Two tailed)} & & \text{(Left tailed)} & & \text{(Right tailed)} \end{array}$$

Step 2 Decide on the significance level, α .

Step 3 Compute the value of the test statistic

$$F = \frac{s_1^2}{s_2^2}$$

and denote that value F_0 .


Procedure 11.3 (cont.)

CRITICAL-VALUE APPROACH

Step 4 The critical value(s) are

$F_{1-\alpha/2}$ and $F_{\alpha/2}$ (Two tailed) or $F_{1-\alpha}$ (Left tailed) or F_{α} (Right tailed)

with $df = (n_1 - 1, n_2 - 1)$. Use Table VIII to find the critical value(s).




Step 5 If the value of the test statistic falls in the rejection region, reject H_0 ; otherwise, do not reject H_0 .

OR

P-VALUE APPROACH

Step 4 The F -statistic has $df = (n_1 - 1, n_2 - 1)$. Obtain the P -value by using technology.



Step 5 If $P \leq \alpha$, reject H_0 ; otherwise, do not reject H_0 .

Step 6 Interpret the results of the hypothesis test.